

Associations between Light Rail Transit and physical activity: a systematic review

Léa Ravensbergen, Rania Wasfi, Mathilde Van Liefferinge, Isidor Ehrlich, Stephanie A. Prince, Gregory Butler, Yan Kestens & Ahmed El-Geneidy

To cite this article: Léa Ravensbergen, Rania Wasfi, Mathilde Van Liefferinge, Isidor Ehrlich, Stephanie A. Prince, Gregory Butler, Yan Kestens & Ahmed El-Geneidy (2023) Associations between Light Rail Transit and physical activity: a systematic review, *Transport Reviews*, 43:2, 234-263, DOI: [10.1080/01441647.2022.2099999](https://doi.org/10.1080/01441647.2022.2099999)

To link to this article: <https://doi.org/10.1080/01441647.2022.2099999>



© 2022 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group



Published online: 12 Jul 2022.



Submit your article to this journal



Article views: 1758



View related articles



View Crossmark data



Citing articles: 1 View citing articles

Associations between Light Rail Transit and physical activity: a systematic review

Léa Ravensbergen  ^a, Rania Wasfi  ^b, Mathilde Van Liefferinge ^a, Isidor Ehrlich ^a,
Stephanie A. Prince  ^{b,c}, Gregory Butler  ^b, Yan Kestens  ^d and
Ahmed El-Geneidy  ^a

^aSchool of Urban Planning, McGill University, Montreal, Canada; ^bCentre for Surveillance and Applied Research, Public Health Agency of Canada, Montreal, Canada; ^cSchool of Epidemiology and Public Health, Faculty of Medicine, University of Ottawa, Montreal, Canada; ^dMontreal University Hospital Research Center, Université de Montréal, Montreal, Canada

ABSTRACT

Investment in public transport is on the rise as many cities around the world aim to reduce their carbon footprint and improve population health. One such investment is building or extending Light Rail Transit (LRT). Focusing on studies in the USA, Canada, Australia, and New Zealand, this paper reports the results of a systematic review on the associations between LRT and physical activity. This systematic review adhered to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement. Twenty studies were identified through a search of five bibliographic databases (Web of Science, Transport Research International Documentation (TRID), Scopus, Medline, and SPORTDiscus) ($n=5,866$) and a systematic Google search ($n=446$). At least two reviewers conducted the search and reviewed the titles and abstract of each identified article to include in the review. Standardized data extraction forms were used to document information from each selected article. The forms included a risk of bias assessment tool. Two reviewers completed the Effective Public Health Practice Project (EPHPP) Quality Assessment Tool for Quantitative Studies. Our findings show that moderate certainty of evidence exists for the relationship between LRT and walking behaviour. Here, all studies, most of which were natural experiments ($n=6$), found a positive association between LRT and walking behaviour, with LRT leading to an increase of 7–40% in walking in most studies ($n=7$ out of 8). A positive relationship between LRT and moderate-to-vigorous intensity physical activity (MVPA) and between LRT and cycling was also often identified; however, results were inconsistent, and certainty of evidence is low for MVPA, and very low for cycling. Further, some studies ($n=3$) identify differences in physical activity participation at different LRT stations, suggesting that station design, surrounding land use, and built environment play important roles in promoting

ARTICLE HISTORY

Received 21 August 2021
Accepted 27 June 2022

KEYWORDS

Light Rail Transit; physical activity; literature review; walking; cycling; MVPA

CONTACT Ahmed El-Geneidy  ahmed.elgeneidy@mcgill.ca  School of Urban Planning, McGill University, Montreal, Canada

© 2022 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group
This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way.

physical activity around LRT. Given this, practitioners can be relatively confident that LRT investments will result in increased walking behaviour.

Introduction

In response to growing environmental (e.g. air quality, climate change) and population health (e.g. physical activity, quality of life, air pollution) concerns, many cities worldwide are increasing their investment in public transport systems. Indeed, public transport has many benefits to both individuals and communities: it can be an affordable travel mode (especially when compared to car use), it can reduce congestion, enhance social connectedness, improve air quality, and increase the quality of life (Sener, Lee, & Elgart, 2016). Unlike driving, public transport also often requires walking to and from stops and is, therefore, expected to encourage physical activity through multimodal travel (Sener et al., 2016). One form of public transport that has become increasingly popular in recent years is Light Rail Transit (LRT) (Sinclair, 2019), defined as fully electric passenger urban rail transit that is partially or fully separated from vehicle traffic (Johnson, 2019; Malouff, 2015; The Transport Politic, 2021). In North America, the number of Light Rail projects under construction grew from 1 in 2012 to 19 in 2019 (Sinclair, 2019). This increase in popularity of LRT may be due to its tendency to have lower capital costs and increased reliability compared to heavy rail systems or that it may encourage transit-oriented development in ways that other, less permanent forms of public transport, such as buses, are unable to do.

Two systematic reviews on public transport and physical activity have been published (Rissel, Curac, Greenaway, & Bauman, 2012; Xiao, Goryakin, & Cecchini, 2019), and one meta-analysis on the effects of rapid transit interventions on physical activity has been conducted (Hirsch, DeVries, Brauer, Frank, & Winters, 2018). Xiao et al. (2019) reviewed the impacts of building, extending, or improving local public transport options on physical activity and found that public transport investments are associated with approximately 30 minutes of additional walking (or other light to moderate physical activity) per week. No significant relationship between new transit and moderate-to-vigorous intensity physical activity (MVPA) was found. Similarly, Rissel et al. (2012) found 8–33 minutes of additional walking was attributable to the use of public transport among adults.

These existing reviews focus on all types of public transport, but LRT may yield different outcomes than other forms of public transport. For instance, riders may be willing to walk further to reach this form of public transport than others (e.g. a bus stop). If secure bicycle parking is available, it is possible that cyclists will travel even greater distances to use a station. Given this and the recent rise in popularity of LRT, this paper presents the results of a systematic review of studies examining the relationship between LRT and physical activity. By focusing on LRT, this paper builds on Hirsch et al. (2018)'s meta-analysis on the effects of rapid transit interventions on physical activity, which found that while transport-related physical activity increased, overall physical activity levels decreased. Their meta-analysis examined LRT in conjunction with Bus Rapid Transit, and Rail Rapid Transit. Five studies were included in this meta-analysis, three of which evaluated the implementation of LRT. Therefore, the objective of this systematic review is to build upon the work of Hirsch et al. (2018), by focusing solely on LRT and expanding their scope by including all

studies, regardless of design, which examined the relationship between this form of public transport and physical activity (e.g. walking, cycling, MVPA). The certainty of evidence for each physical activity outcome (e.g. walking, cycling, MVPA) was also assessed to provide policy makers with a clear indication of how confident they can be that LRT investments will result in different physical activity outcomes. Because research on LRT and physical activity originating from places with high public transport and active travel rates, such as many European and Asian cities, might find stronger evidence than that of cities with more auto-oriented urban planning legacies, the geographic scope of this paper is focused on LRT in the United States, Canada, Australia, and New Zealand to ensure policy makers in these contexts do not over-estimate the effect of this public transit investment. Further, this reduced scope allows for a rigorous exploration of the risk of bias in each paper and the certainty of evidence of the body of work. The secondary objective of our study is to document the different approaches used in this body of work. We do so by documenting the theoretical frameworks guiding the included studies, as well as whether previous research considered equity or self-selection, two important topics in research on travel behaviour.

Methods

Context and study inclusion and exclusion criteria

This systematic review adhered to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement. A protocol for this review was prospectively registered on Open Science Framework (<https://osf.io/ux5vn/>) and PROSPERO (registration number: CRD42021254690).

Population: No age restrictions were placed on the participants in the studies. Only studies that evaluated the association between LRT and physical activity in the United States, Canada, Australia, and New Zealand were eligible. These countries were selected because their cities tend to have similar urban fabric and built environments and previous research has found similar factors influence walking to public transport in these countries, as opposed to differences in Asia and Europe (van Soest, Tight, & Rogers, 2020).

Exposure: Following guidelines put forth by Malouff (2015) and Johnson (2019), and Light Rail data provided by The Transport Politic (2021), LRT was defined as fully electric passenger urban rail transit that is partially or fully separated from vehicle traffic. Studies that concentrated on streetcar or street railroads, bus rapid transit or rail rapid transit were excluded from this systematic review. LRT could be assessed through self-reported use, objectively measured use, or proximity to LRT (e.g. living near a station). Studies that considered LRT use alongside other forms of public transport (i.e. did not separate the modes in the analysis) were excluded (Bopp, Gayah, & Campbell, 2015; Carlson, Watson, Paul, Schmid, & Fulton, 2017; Lachapelle & Noland, 2012; Lachapelle & Pinto, 2016; Langlois, Wasfi, Ross, & El-Geneidy, 2016).

Outcomes: The primary outcome was physical activity. Physical activity could be device-measured (e.g. pedometer, accelerometer), or self-reported (e.g. questionnaire with reported minutes of walking or biking or a number of times engaged in an activity) and did not have to be a direct result of using an LRT. Papers that examined the impact of LRT on BMI or body weight that did not look at physical activity separately (e.g. Brown, Smith, Jensen, & Tharp, 2017) were excluded as body weight is a complex outcome

influenced by myriad factors that extend beyond physical activity. Studies of simulations (e.g. anticipated physical activity if LRT investment was made) were excluded, as they do not present real-world physical activity data. Secondary outcomes included documentation of the theoretical frameworks guiding the research, and whether the research considered equity and participant self-selection.

Study designs: All original studies that empirically examined the associations between LRT and physical activity in the selected countries were included, regardless of the study design.

Publication status and language: Based on the authors' language capabilities, only publications in English or French were eligible. Abstracts without full texts were excluded. When two papers reported on the same data, peer-reviewed publications were prioritized (e.g. in the case of a dissertation and a peer-reviewed publication, only the peer-reviewed publication was retained), as well as the most recent source (e.g. newer publications prioritized before older publications).

Search strategy

The search strategy was developed by the authorship team in consultation with a librarian specializing in systematic reviews. Two search strategies were developed: one for academic literature and one for grey literature. Searches were conducted in May 2021.

The academic literature search included five electronic bibliographic databases: Web of Science, Transport Research International Documentation (TRID), Scopus, Medline, and SPORTDiscus. Trial searches were run and expanded upon to ensure that pre-identified key papers were captured by the search strategy. Titles, abstracts, and keywords (when possible, TRID only allowed a title search) were searched for synonyms of LRT and physical activity. The full electronic search strategy, including all search terms, can be found in Appendix 1.

The search of academic articles was complimented by a search of the grey literature. Firstly, the ProQuest Dissertations & Theses Global database, the world's most comprehensive collection of dissertations and theses, was searched. The same search terms used for the journal databases were used to search titles, abstracts, and keywords.

Google searches were used to search for each city known to have LRT based on a database from The Transport Politic (2021) (Appendix 2) by searching for the city's name alongside the terms "light rail transit" and "physical activity". This was done on a browser with no previous history in incognito mode. The two reviewers reviewed the first two pages of the resulting search and compiled all potentially relevant reports. Eligible reports needed to present data on the associations between LRTs and physical activity that were not already presented in the academic literature or dissertation search. If a report was selected for inclusion, they continued their search to the following page. This was done until a full page with no selected reports were identified. Though the aim was to be comprehensive, it is, of course, possible that reports further down the search were missed. This is more likely for reports that do not include the key terms (e.g. physical activity, LRT) in locations that are picked up easily by search engines (e.g. title, keywords, etc.). The two reviewers then met to compare the resulting reports. On the rare occasion that the two reviewers did not identify the same reports through the Google search, they resolved which to include through discussion. For example, this

occurred when a grey literature document identified by one reviewer (and not the other) presented data that was already presented in an academic publication.

Finally, the reference lists of all included studies, as well as all literature review papers identified through this screening process, were hand-searched to identify potentially eligible articles.

Selection of studies

For the academic literature, abstract and full-text screening was conducted by two independent reviewers using Rayyan software, an open-source systematic review manager. If conflicts arose, they were discussed with a third reviewer; the resulting articles' full texts were screened by both reviewers for inclusion. When conflicts arose, they were resolved through discussion. For the grey literature search, two reviewers independently examined the first two pages of the resulting search and compiled all potentially relevant reports. If a report was selected for inclusion, they continued their search to the following page. This was done until a full page with no selected reports were identified. The two reviewers then met to compare the resulting reports selected and to resolve conflicts through discussion. Eligible reports needed to present data that were not already present in the peer-reviewed published literature or dissertation search.

Data extraction

Standardized data abstraction forms were independently completed by two reviewers using Google Forms. A Google Form was developed with detailed instructions to extract information from studies to minimize potential bias in extraction between reviewers. Extracted data included: authors, title, publication year, study setting (country and city), study context, seasonality of data collection, study design, population description, research question, LRT exposure measure, how exposure was defined, physical activity outcome measure, whether physical activity measure was self-reported or device-measured, confounders measured, theoretical framework, statistical methods used, results, whether the study considered equity in any way (e.g. gender, income, socio-economic status (SES) indicators, or race/ethnicity-based differences in physical activity or LRT exposure), future research recommendations, and policy recommendations. Once both reviewers completed data extraction, the extractors met to review and validate the data. Any disagreements that arose were resolved through discussion.

Risk of bias

The standardized data extraction forms included a risk of bias assessment. Both reviewers completed the Effective Public Health Practice Project (EPHPP) Quality Assessment Tool for Quantitative Studies (<https://www.epph.ca/quality-assessment-tool-for-quantitative-studies/>). The instrument assesses eight domains: (1) selection bias; (2) study design; (3) confounders; (4) blinding; (5) data collection method; (6) withdrawals/dropouts; (7) intervention integrity; and, (8) analyses. Reviewers rated each domain as strong, moderate, or weak. The tool was modified to better reflect the topic of this review and guidelines were established to help the reviewers give the

papers consistent grades (Appendix 3). Once complete, the two reviewers compared their risk of bias assessments. If disagreements arose, they were resolved through discussion. Each study received a global rating based on EPHPP criteria (i.e. "strong" if no domains had a weak rating, "moderate" if one domain had a weak rating, and "weak" if two or more domains had a weak rating).

Evaluation of certainty of evidence

The certainty of evidence was then assessed using a modified GRADE (Grading of Recommendations Assessment, Development, and Evaluation) approach (<https://gdt.gradepro.org/app/handbook/handbook.html>). Certainty of evidence was assessed as "high", "moderate", "low", or "very low" for each outcome (e.g. walking behaviour, cycling behaviour, MVPA).

Each outcome was assigned an initial certainty of evidence based on the study design. If the designs were randomized controlled trials/natural experiment/quasi-experimental




Figure 1. PRISMA flow diagram.

studies, the evidence was assigned an initial level of “high”, whereas cross-sectional studies were assigned an initial level of “low”. The certainty of evidence for each outcome was assessed separately for the experimental and observational evidence. The final certainty of evidence was based on the evidence from the highest quality study design available (e.g. experimental above observational).

The certainty of evidence was based on the following: risk of bias, inconsistency, indirectness, imprecision, or publication bias. In instances where the certainty of evidence was not downgraded, it could be upgraded if the following were observed: large effect, dose-response, or opposing bias and confounders (Balshem et al., 2011; Murad, Mustafa, Schünemann, Sultan, & Santesso, 2017). The specific evaluation criteria guiding the evaluation of evidence can be found in Appendix 4.¹ One reviewer independently assessed the certainty of evidence, and a second reviewer verified the assessments.

Results

The database search identified a total of 5866 articles, 1969 of which were duplicates. Title and abstract screening resulted in 38 potential articles whose full texts were screened by both reviewers for inclusion. Many articles were removed due to their lack of focus on LRT, their combined focus on LRT with other forms of public transport, their lack of focus on measured physical activity, or their geographic focus. After full-text screening, 12 papers were included in this review. The grey literature search identified 440 dissertations and theses, 3 of which were included, and 5 eligible reports from the Google search (Figure 1).

Summary of results

A summary of key characteristics of the papers, dissertations, and reports included in this review can be found in Table 1 ($n = 20$). Most of the studies ($n = 17$) took place in the United States, including all 12 academic articles. The most frequent study setting was Salt Lake City ($n = 5$).

Primary objective: LRT and physical activity

Eleven studies used a natural experiment research design (before and after), while nine were cross-sectional (Table 1). The natural experiment studies all examined physical activity before and after the introduction of a new LRT or LRT extension. Spears et al. (2017) was the only study to examine these effects at multiple time periods after opening (6 and 18 months after opening). Three of the natural experiments examined the effects of LRT on physical activity at different distances to the station (Brown, Smith, et al., 2016; Hong et al., 2016; Huang et al., 2017), while two compared the area around the LRT to a control group (Ewing & Hamidi, 2014; Spears et al., 2017). For instance, Spears et al. (2017) compared physical activity amongst residents living within walking distance of new LRT stations to that of residents living in a neighbourhood with a similar built environment and socio-demographic characteristics further from the station. The use of comparative groups was common among the cross-sectional studies. The most common comparative analyses were by distance to LRT (Kumar

Table 1. Key characteristics of included studies.

Paper	LRT location	Sample size analysed	Comparative group	Measurement tools/self-reported or measured	LRT exposure measure	Physical activity measure
Natural experiments						
Barbaric and Alizadeh (2017)	Australia, Gold Coast	NR	Before and after (time period NR)	Measurement tools not reported/NR	Proximity to LRT: NR	Walking behaviour: pedestrian counts Cycling behaviour: cyclist counts
MacDonald, Stokes, Cohen, Kofner, and Ridgeway (2010)	USA, Charlotte	498	Before and after (~1 year)	Survey/self-reported	LRT use: intention to use LRT, and LRT use during daily commute	MVPA: modified version of the International Physical Activity Questionnaire RPA: vigorous activity 3 times per week (20 minutes or more per time) or walking 5 times per week (30 minutes or more per time)
Hong, Boarnet, and Houston (2016)	USA, Los Angeles	204 and 73 accelerometer sample	Before and after (~1 year) Near and far	MVPA: accelerometry/measured Walking behaviour: travel survey/self-reported (validated against GPS data)	Proximity to LRT: residing <½ mile and >½ mile from a new LRT station	MVPA: average daily minutes of MVPA Walking behaviour: average daily walking trips
Spears, Boarnet, and Houston (2017)	USA, Los Angeles	285 6 months before, 208 6 months after, and 173 18 months after	Before and after (6 and 18 months) control vs. experimental	Seven-day travel log/ self-reported	Proximity to LRT: residences within 1 km vs. 1–3 km of LRT LRT use: trip frequency	Walking behaviour: frequency walking trips Cycling behaviour: frequency bicycle trips
Hampton Roads Transit (2015)	USA, Norfolk, Virginia	NR	Before and after (2 years)	Travel survey/self-reported	LRT use	Walking and cycling behaviour: access to LRT using non-motorized mode (walk or bike)
Ewing and Hamidi (2014)	USA, Portland	65	Before and after (17 years) Near LRT vs. control group	Walking and cycling behaviour: travel diary/ self-reported	Proximity to LRT: compared behaviour of those living within an LRT corridor and those in a control corridor	Walking behaviour: walking trips by purpose and by location Cycling behaviour: bike trips by purpose and by location
Brown and Werner (2007)	USA, Salt Lake City	51	Before and after (one year)	Travel survey/self-reported Accelerometry/ measured	Frequency of LRT use	MVPA: bouts of moderate activity (at least 1952 counts per minute)
Brown, Smith, et al. (2016)	USA, Salt Lake City	536 (910 first wave)		Accelerometry and GPS/measured	Proximity to LRT station: living near (<800 m) and	

(Continued)

**Table 1.** Continued.

Paper	LRT location	Sample size analysed	Comparative group	Measurement tools/self-reported or measured	LRT exposure measure	Physical activity measure
Brown, Werner, et al. (2016)	USA, Salt Lake City	536	Before and after (one year) Near vs. far Four transit ridership groups	GPS and accelerometer/ measured	far (801–2000 m) with tests for alternate distances (<600 and <1000 m) Identify residents who made LRT trips in the complete street corridor	Walking behaviour: walking trips, transit-related active travel trips Cycling behaviour: trip frequency MVPA: minutes (threshold set at 2020 CPM)
Miller et al. (2015)	USA, Salt Lake City	536	Before and after (~1 year)	GPS and accelerometer/ measured	LRT use: participants categorized as never, continued, new, or former transit user	LMPA: bout with a minimum duration of 5 minutes with a minimum of 1000 accelerometer counts per minute (cpm)
Huang, Moudon, Zhou, Stewart, and Saelens (2017)	USA, Seattle	198	Before and after (~2 years) Near and far	PA: accelerometer and GPS/measured LRT: travel survey/self-reported	LRT use: trip frequency Distance to LRT: live within 1 mile	Walking behaviour: walk trip frequency within ¼ mile of stations
Cross-sectional studies						
Eady and Burtt (2019)	Australia, Melbourne	NR	Distance walked	Measurement tools NR	LRT use	Walking distance: distance
O'Sullivan (1995)	Canada, Calgary	1843	Different stations	Survey of walking routes (self-reported) of people at LRT stations (measured)	LRT use: surveyed people at LRT stations	Walking distance: survey of walking routes
Appleyard, Frost, and Allen (2019)	USA, Charlotte, Dallas, Denver, Houston, Minneapolis-St. Paul, Phoenix, Portland, Sacramento, Salt Lake, San Diego, San Jose, and Seattle	NA – population-level variables across 357 stations in 12 metro areas	Categorizes and compares stations based on an array of variables	Travel survey/self-reported	Proximity to LRT: half mile straight-line distance to light rail transit	Walking behaviour: walk to work Cycling behaviour: bike to work
Hess, Ray, and Attard (2014)	USA, Buffalo	1397	Free pass holders ($n = 643$) vs. control group (754)	Travel survey/self-reported	LRT use and behaviour during a pilot transit pass programme	Walking and cycling behaviour: frequency and minutes walking or cycling during commute
Kumar Maghelal (2007)	USA, Dallas	1022	Built environment at stations near vs. far	Travel survey/self-reported	Proximity to LRT: ¼ and ½ mile from stations	Walking behaviour: % of LRT passengers who walk to LRT
Schoner and Cao (2014)	USA, Minneapolis	1303	LRT corridors vs. control areas	Travel survey/self-reported	LRT proximity: LRT corridor vs control corridors	Walking behaviour: utilitarian walking (shopping trips) and recreational walking (strolling)

Noland, Ozbay, DiPetrillo, and Iyer (2014)	USA, New Jersey	1629	Near vs. far	Survey/self-reported	LRT proximity: residence near LRT (within ½ mile) vs. far (½ mile – 2 miles) Also asked about LRT use, but not in relation to PA	Walking behaviour: multiple types of walking (walking frequency, outdoor walking frequency, commute mode, walk to restaurant or coffee shop, etc.) Cycling behaviour: bicycle as access mode to transit
Crist et al. (2021)	USA, San Diego	6894	None	Travel survey/self-reported	Anticipated use of LRT and distance to LRT	Walking and cycling behaviour: odds of and duration of transport-related physical activity
McAslan (2018)	USA, Seattle	249 surveys and 43 interviews	Not in relation to physical activity	Interviews/self-reported	LRT use	Walking distance

Note: LRT = Light Rail Transit, LMPA = light-to-moderate intensity physical activity, MVPA = moderate-to-vigorous intensity physical activity, NR = not reported, RPA = Recommended Physical Activity, USA = United States of America.

Table 2. Summary of key findings.

Outcome		Study	Key findings
Walking	Natural experiment	Brown, Smith, et al. (2016)	<ul style="list-style-type: none"> 38% (from 34% to 47%) and 64% (from 11% to 18%) increase in residents completing non-transit walking trips near and far (respectively, specific distance not stated) from LRT after operational Odds of non-transit walks for those living near (<800 m) and far (1000 m) from LRT post-construction is 0.28 (95% CI: 0.19–0.44)
		Ewing and Hamidi (2014)	<ul style="list-style-type: none"> Daily walking trips (all purposes) increased in LRT corridor after construction from 0.86 trips to 2.16 (an increase of 151%) ($p < .001$) More weekly walk trips in the LRT corridor after construction than control corridor ($p = .018$)
		Hong et al. (2016)	<ul style="list-style-type: none"> Living within a half mile buffer of LRT positively associated with self-reported walk trip counts for all purposes Average daily walking trips increased from 1.00 to 1.29 after LRT operational (an increase of 29%) ($p < .05$) Those living in control area exhibited a lower number of walking trips (average daily trips = 0.79). Proportion unchanged after LRT operational
		Spears et al. (2017)	<ul style="list-style-type: none"> Six months after LRT stop operational, average daily all-purpose walking trips increased from 1.58 to 1.75 (a 10.76% increase) ($p < .05$) 18 months after LRT operational, no statistically significant change identified (mean number returned to 1.58)
		Huang et al. (2017)	<ul style="list-style-type: none"> Minutes walking decreases 36.2–25.1 ($p < .001$) after LRT Walking within one quarter Euclidean mile of an LRT station increased from 7.7 to 8.4 average daily minutes ($p = .413$) Per cent station-area walking increased from 0.20 to 0.26, a 30% increase, $p < .001$
		Barbaric and Alizadeh (2017)	<ul style="list-style-type: none"> Pedestrian counts generally remained stable, however, they increased on light rail pedestrian links
	Cross-sectional	Schoner and Cao (2014)	<ul style="list-style-type: none"> Travel attitudes, residential preferences, and built environment characteristics associated with walking ($p < .1$) Living in LRT area (quarter mile buffer) not statistically significantly associated with walking to the store ($p > .1$) Mean walking trips for leisure and to the store consistently higher in LRT corridor (3.68 mean strolling trips and 2.04 mean walking to the store trips) than the two suburban controls (3.22 and 3.52 (strolls) and 1.88 and 1.93 (store)), but not consistently higher than the two urban controls (3.67 and 3.51 (strolls) and 1.96 and 2.35 (store))
		Noland et al. (2014)	<ul style="list-style-type: none"> Walking frequency, walking as a commute mode in the last week, walking as an access mode for transit, and per cent of walking to other destinations (restaurant or coffee shop, food or grocery shopping) higher amongst those living near the station (within $\frac{1}{2}$ mile) than those living further away ($\frac{1}{2}$ mile – 2 miles). Frequency of walking outdoors for five or more minutes was higher amongst those living near LRT (within $\frac{1}{2}$ mile) than farther ($\frac{1}{2}$ mile – 2 miles) (58.2% vs. 41.8%) Differences not statistically tested

(Continued)

Table 2. Continued.

Outcome		Study	Key findings
Cycling	Natural experiment	Brown, Smith, et al. (2016)	<ul style="list-style-type: none"> • 2% increase in cycling near LRT after construction • Relationship only statistically significant when comparing to residents living far from the LRT <i>and</i> before construction ($p = .04$)
		Ewing and Hamidi (2014)	<ul style="list-style-type: none"> • Bicycle trips decreased from 0.12 to 0.11 ($p = .88$) after LRT construction • Bicycle trips only marginally greater in LRT corridor than control corridor (0.24 vs. 0.11, $p < .1$)
		Spears et al. (2017)	<ul style="list-style-type: none"> • Mean bicycle trips went from 0.17 before to 0.34 at 6 months and 0.24 at 18 months • LRT users complete more bicycle trips than non-users both before the LRT was built and at two time points afterwards (6 and 18 months)
		Barbaric and Alizadeh (2017)	<ul style="list-style-type: none"> • Cyclist-counts increased after LRT construction (extent and statistical significance not reported)
	Cross sectional	Noland et al. (2014)	<ul style="list-style-type: none"> • Bicycle access mode for transit more common amongst those living near LRT than those living farther (33.3% vs. 66.7%, statistical significance not reported)
Walking and cycling	Cross sectional	Crist et al. (2021)	<ul style="list-style-type: none"> • Positive relationship between distance to LRT station and both the odds of participating in and duration of transport-related physical activity ($p > .1$)
		Hess et al. (2014)	<ul style="list-style-type: none"> • During free transit pass programme, 58% of respondents reported walking or cycling more during their commute • Of those, 18% reported an increase in their walking or cycling by 11–15 minutes per day and 19% reported an increase of 16–20 minutes per day
	Natural experiment	Hampton Roads Transit (2015)	<ul style="list-style-type: none"> • 73% of riders travelled to the transit stop by foot or bicycle
	Cross sectional	Eady and Burtt (2019)	<ul style="list-style-type: none"> • Residents walked a mean distance of 390 m to reach a bus stop, and 360 m to reach an LRT stop
		McAslan (2018)	<ul style="list-style-type: none"> • Respondents stated they would walk farther to reach an LRT stop than a bus stop
		O'Sullivan (1995)	<ul style="list-style-type: none"> • Residents willing to walk further to reach an LRT stop (422.3 m) compared to the City's walking distance guidelines (400 m) • Residents who walk frequently to or from LRT stations walk further than those who do not
MVPA	Natural experiment	Brown and Werner (2007)	<ul style="list-style-type: none"> • Mean bouts of moderate-intensity physical activity did not change (0.06 at before and after) after LRT • Self-reported rail ridership related to bouts of moderate activity both before and after LRT construction
		Brown, Werner, et al. (2016)	<ul style="list-style-type: none"> • MVPA significantly higher amongst continuing riders (33.62 minutes of MVPA per 10 hours), former riders (28.78), and new riders (23.84) compared to never riders (17.33)
	Hong et al. (2016)		<ul style="list-style-type: none"> • Average daily minutes of MVPA decreased from 23.09 minutes before LRT to 21.52 after LRT ($p > .05$)

(Continued)

Table 2. Continued.

Outcome	Study	Key findings
	MacDonald et al. (2010)	<ul style="list-style-type: none"> Odds ratio of increasing PA through vigorous exercise to meet RPA levels associated with LRT use (OR: 3.32, 95% CI 0.81, 3.63), but only significant at $p < .10$
Odds of meeting RPA	Natural experiment MacDonald et al. (2010)	<ul style="list-style-type: none"> Odds of meeting recommended physical activity levels (vigorous activity 3 times a week, ≥ 20 minutes a time; or walking 5 times a week, ≥ 30 minutes a time) higher with LRT use, but not statistically significant
LMPA	Natural experiment Miller et al. (2015)	<ul style="list-style-type: none"> LMPA (total and in relation to transit use) decreased amongst those that discontinued using transit and increased amongst those who began using transit (from 22.03 to 27.30 average minutes) Relationship between LRT use and total physical activity not statistically significant, but the relationship with transit-related physical activity was ($p < .001$)
Physical activity across LRT	Cross-sectional Appleyard et al. (2019)	<ul style="list-style-type: none"> 18.05% and 3.83% walked to coordinating and emerging stations, respectively 4.35% and 1.71% cycled at coordinating and emerging stations, respectively
	Kumar Maghelal (2007)	<ul style="list-style-type: none"> No significant relationship between built environment variables and walking to transit identified at half a mile from the station Density was found to explain walking to the station at one-quarter of a mile ($p < .05$)
	O'Sullivan (1995)	<ul style="list-style-type: none"> People walked 651.4 and 326.4 m, on average, to reach suburban and downtown stations, respectively

Note: LRT = Light Rail Transit, LMPA = light-to-moderate intensity physical activity, MVPA = moderate-to-vigorous intensity physical activity, RPA = recommended physical activity.

Maghelal, 2007; Noland et al., 2014) and across stations or station typologies (Appleyard et al., 2019; Kumar Maghelal, 2007).

Nine studies utilized proximity to LRT as the exposure measure, while ten looked at LRT use and two studied both (Table 1). The LRT exposure was self-reported in nine studies and measured in 10 (2 did not specify). The physical activity measures included walking behaviour, walking distance, cycling behaviour, walking and cycling behaviour combined, MVPA, light-to-moderate intensity physical activity (LMPA), and meeting recommended levels of physical activity (e.g. ≥ 150 minutes per week). Other studies compared physical activity across LRT stations.

A summary of key findings can be found in Table 2. The quality of the studies was evaluated using the EPHPP Quality Assessment Tool (Table 3). Using the EPHPP tool, two peer-reviewed papers received a “weak” global rating, while seven of the eight grey literature documents received this “weak” rating. Two best practices that were frequently omitted from the methods sections of these papers included comparing the study sample to the general population (e.g. by comparing the sample’s characteristics to the census) and reporting the number and reasons for withdraws or dropouts in the natural experiment studies.

The overall certainty of the evidence for each physical activity outcome discussed in the papers was assessed using the GRADE Evidence Profile (Table 4).

In the following section, we provide a narrative synthesis of the study's results organized by the type of physical activity under study. An analysis of these findings, including how best researchers can respond to remaining knowledge gaps, follows in the Discussion section of the paper.

Walking behaviour

The most studied physical activity outcome was walking behaviour (Table 2). Of the eight studies focused on the association between LRT and walking behaviour, six were natural experiments, all of which found evidence for a positive association between walking and LRT. Four of these six studies found statistically significant evidence between the construction of LRT and increased walking behaviour (Brown, Smith, et al., 2016; Ewing & Hamidi, 2014; Hong et al., 2016; Spears et al., 2017). Spears et al. (2017) was the only study that looked at the physical activity effects of light rail at two follow-up times. They found that walking trips increased amongst residents in neighbourhoods with an LRT stop six months after opening but that no statistically significant change was identified after 18 months. Amongst the two that did not find consistent statistically significant evidence, one was a grey literature report that did not indicate how walking was measured nor the extent to which it increased (Barbaric & Alizadeh, 2017). In the other, walking behaviour decreased after the LRT opened ($p < .001$), however per cent station-area walking increased by 30%. In the two cross-sectional studies, Schoner and Cao (2014) found mixed evidence and Noland et al. (2014) found consistent evidence, but results were not statistically tested (Table 2).

For walking behaviour, there was moderate certainty of evidence for a positive association with LRT. Across studies that reported quantitative results, LRT was generally associated with a 7–40% increase in walking, with one study reporting a 151% increase.

Cycling behaviour

The relationship between LRT and cycling behaviour was less extensively studied compared to walking behaviour ($n = 5$). The results are also less consistent amongst both natural experiments and cross-sectional studies (Table 2). Many studies found the expected relationship, but either only found statistical significance in certain scenarios (Brown, Smith, et al., 2016; Ewing & Hamidi, 2014), found no significance (Spears et al., 2017), or did not check for significance (Barbaric & Alizadeh, 2017; Noland et al., 2014). The overall certainty of evidence for cycling behaviour is "very low" (Table 3).

Walking and cycling behaviour

Three papers combined walking and cycling behaviour (Table 2). All present findings from self-reported behaviour in travel surveys, and all find evidence for a relationship between LRT and physical activity, but its statistical significance was either low or not reported. Crist et al. (2021) is the only peer-reviewed article focused on walking and cycling combined, and it focuses on a University setting. Two reports also exist, one examining the impacts of a free transit pass pilot programme in Buffalo, New York (Hess et al., 2014), the other examining the impacts of the LRT in Norfolk, Virginia (Hampton Roads Transit, 2015).

Overall, there was "very low" certainty of evidence for a positive association between LRT and walking and cycling behaviour (Table 3).

**Table 3.** Risk of bias assessment.

Paper	Global rating	Low selection bias	Study design	Confounder	Blinding	Data collection methods	Withdrawals/ dropouts	Intervention integrity	Analyses
Peer-reviewed journal articles									
Appleyard et al. (2019)	Weak	Moderate	Weak	Moderate	Strong	Weak	NA	NA	Moderate
Brown, Smith, et al. (2016)	Strong	Moderate	Strong	Strong	Moderate	Strong	Strong	Strong	Moderate
Brown, Werner, et al. (2016)	Strong	Moderate	Strong	Moderate	Moderate	Strong	Strong	Strong	Strong
Brown and Werner (2007)	Moderate	Weak	Strong	Moderate	Moderate	Strong	Moderate	Strong	Strong
Crist et al. (2021)	Moderate	Strong	Weak	Moderate	Moderate	Moderate	NA	NA	Strong
Ewing and Hamidi (2014)	Weak	Strong	Strong	Moderate	Moderate	Weak	Weak	Weak	Moderate
Hong et al. (2016)	Strong	Moderate	Strong	Strong	Moderate	Strong	Moderate	Moderate	Strong
Huang et al. (2017)	Strong	Moderate	Strong	Strong	Moderate	Strong	Moderate	Strong	Strong
MacDonald et al. (2010)	Moderate	Moderate	Strong	Moderate	Moderate	Weak	Strong	Moderate	Moderate
Miller et al. (2015)	Strong	Moderate	Strong	Moderate	Moderate	Strong	Moderate	Moderate	Moderate
Schoner and Cao (2014)	Strong	Strong	Moderate	Moderate	Moderate	Moderate	NA	NA	Moderate
Spears et al. (2017)	Strong	Strong	Strong	Strong	Strong	Moderate	Strong	Moderate	Moderate
Grey literature									
Barbaric and Alizadeh (2017)	Weak	Weak	Weak	Weak	Moderate	Weak	Weak	Weak	Weak
Eady and Burtt (2019)	Weak	Weak	Weak	Weak	Moderate	Moderate	NA	NA	Weak
Hampton Roads Transit (2015)	Weak	Weak	Strong	Weak	Moderate	Weak	Weak	Moderate	Weak
Hess et al. (2014)	Weak	Weak	Weak	Moderate	Moderate	Moderate	NA	NA	Weak
Kumar Maghel (2007)	Weak	Strong	Weak	Weak	Strong	Moderate	NA	NA	Strong
McAslan (2018)	Weak	Weak	Weak	Weak	Moderate	Weak	NA	NA	Weak
Noland et al. (2014)	Strong	Moderate	Moderate	Moderate	Moderate	Moderate	NA	NA	Strong
O'Sullivan (1995)	Weak	Moderate	Weak	Weak	Moderate	Weak	NA	NA	Moderate

Table 4. GRADE evidence profile.

Outcomes across studies	Study design	Number of studies	Number of participants ^a	Relative effect	Quality of evidence (GRADE)
Walking behaviour	Natural experiment	6	1176	Positive and statistically significant association between LRT and walking behaviour found in most (4/6) studies	Moderate  Initial rating: high Risk of bias: 0 Inconsistency: 0 Indirectness: 0 Imprecision: 0 Publication bias: -1
	Cross-sectional	2	2932	Positive association between LRT and walking behaviour found in one paper, inconsistent evidence found in the other	Very low  Initial rating: low Risk of bias: 0 Inconsistency: -1 Indirectness: 0 Imprecision: -1 Publication bias: 0
Cycling behaviour	Natural Experiment	4	774	Positive association identified in most studies (3/4), but most results not statistically significant	Very low  Initial rating: high Risk of bias: -0.5 Inconsistency: 0 Indirectness: 0 Imprecision: -1 Publication bias: -1
	Cross-sectional	1	1629	Positive association identified, but statistical significance not reported	Very low  Initial rating: low Risk of bias: 0 Inconsistency: NA Indirectness: 0 Imprecision: -1 Publication bias: -1
Walking and cycling behaviour	Natural experiment	1	NR	Positive association identified, but statistical significance not reported	Very low  Initial rating: high Risk of bias: NA Inconsistency: NA Indirectness: -1 Imprecision: -1 Publication bias: 0
	Cross-sectional	2	7537	Positive association identified, but results are not statistically significant in one study, and not reported in the other	Very low  Initial rating: low Risk of bias: 0 Inconsistency: 0 Indirectness: -1

**Table 4.** Continued.

Outcomes across studies	Study design	Number of studies	Number of participants ^a	Relative effect	Quality of evidence (GRADE)
MVPA	Natural experiment	4	1158	2/4 studies find significant positive association. One study finds no association and the other finds negative, but statistically insignificant, association	Low
LMPA	Natural experiment	1	536	Positive association identified between LRT and total LMPA (not significant) and transit related LMPA (significant)	Moderate
Recommended physical activity	Natural experiment	1	498	Positive, but not significant, association identified	Moderate
Walking distance to LRT stations	Cross-sectional	3	2092	2/3 studies find that respondents walk further to reach LRT stops than other public transit	Very low
Physical activity across stations	Cross-sectional	3	2865	All studies find some variation in physical activity across LRT stations	Very low

Note: LRT = Light Rail Transit, MVPA = moderate-to-vigorous intensity physical activity.

^aWhen multiple samples were reported, most relevant or smallest included herein.

Walking distance

A further three studies examined the distance walked to LRT ([Table 2](#)). An older study set in Calgary, Canada, found that those who walk frequently to or from LRT stations walk further than those who do not. This result points to the variation in walking distances across people using the same LRT system. This study and a more recent study in Seattle, USA, found that people are willing to walk further to reach an LRT stop than a bus stop ([McAslan, 2018](#)). The third study, however, found that residents in Melbourne, Australia walked but less far to reach an LRT stop than a bus stop ([Eady & Burtt, 2019](#)).

Based on the GRADE evaluation, there is very low certainty of evidence for a positive association between LRT and walking distance, whereby individuals are more likely to walk further to access LRT than other forms of public transport.

MVPA

Four natural experiments examined MVPA, three of which measured this outcome through accelerometry ([Brown, Werner, et al., 2016](#); [Brown & Werner, 2007](#); [Hong et al., 2016](#)) ([Table 2](#)). In half of these studies, MVPA either decreased ([Hong et al., 2016](#)), or did not change ([Brown & Werner, 2007](#)) after an LRT was introduced. However, in [Brown, Werner, et al. \(2016\)](#), MVPA was significantly higher amongst all categories of LRT riders than never riders and self-reported MVPA increased with the introduction of LRT ($p < .1$) in [MacDonald et al. \(2010\)](#). Further, in both [Brown and Werner \(2007\)](#) and [Hong et al. \(2016\)](#), MVPA was higher amongst those living near LRT both before construction and after, suggesting the potential for residual confounding due to self-selection. [Noland et al. \(2014\)](#) was the only cross-sectional study that examined MVPA; however, it was only analysed in relation to self-reported health and not physical activity.

There is low certainty of evidence for the relationship between LRT and MVPA. Though half the studies (2/4) find a significant positive association, the others either find no relationship or an insignificant negative association.

Other physical activity measures

Two additional physical activity outcomes were examined. Moderate certainty of evidence exists for each; however, each was only measured in one study. [MacDonald et al. \(2010\)](#) identified a positive, but not statistically significant, relationship between LRT use and the odds of meeting recommended physical activity levels. [Miller et al. \(2015\)](#)'s study in Salt Lake City also found the expected relationship between LRT and physical activity, this time LMPA measuring. LMPA generally encompasses walking; therefore, these results may be used to support the evidence on the associations between LRT and walking behaviour.

Physical activity across stations or station typologies

Three studies compared walking or cycling across LRT stations rather than measuring the associations between LRT and these behaviours. [Appleyard et al. \(2019\)](#) categorized LRT stations in 12 metro areas as emerging (infrequent transit, limited transport connectivity, and segregated/low-intensity land uses), transitioning (moderate transit frequency,

moderate street connectivity, moderate-to-high intensities, and some mixes of uses), or coordinating (high transit frequency, high connectivity, moderate-to-high intensity, and a mix of complementary uses). Here, both self-reported walking and cycling were significantly higher at coordinating stations than emerging stations. Further, density was found to explain walking to the station in one study (Kumar Maghelal, 2007), while in another people walked farther to reach suburban stations than downtown stations (O'Sullivan, 1995). Only three studies compared stations, and the certainty of evidence is very low, however, this research highlights the importance of considering station design when measuring the effects of LRT on physical activity.

Secondary objectives: theoretical frameworks, equity, and self-selection

Most studies did not specify a theoretical framework. Of those that did, the most common was the socio-ecological model, where physical activity is understood as the outcome of dynamic interrelations among personal and environmental factors at different scales (Brown, Smith, et al., 2016; Brown, Werner, et al., 2016; Crist et al., 2021). Appleyard et al. (2019) drew on Smart Growth and livability principles to develop an evaluation typology. McAslan (2018) framed their work on the theory of urban fabrics. This theory highlights how cities comprise multiple urban fabrics based on travel modes (walking, transit, car, etc.), that each requires its own planning approach to alter the automobile urban fabric to improve transit accessibility and walkability.

Few studies explicitly examined whether the physical activity outcomes of LRT were distributed equitably amongst the population, though most included socio-demographic variables in their models (e.g. sex, income, race/ethnicity, education, etc.). For instance, after controlling for preferences, perceptions, and the built environment, Schoner and Cao (2014) found that having a lower income was associated with utilitarian pedestrian trips. One exception is Appleyard et al. (2019), who examined how measures of social vulnerability and exclusion (unemployment, education, poverty, linguistic isolation, and race/ethnicity) varied between different station typologies in California. Results indicate that stations with the highest transit frequency, connectivity, and intensity of land uses had lower social vulnerability and exclusion. These stations were also associated with higher levels of walking and cycling. Further, current studies focus on adult populations. In cases where data on youth or older adults was collected, it was not presented separately from that of adults.

Few articles explicitly discussed residential self-selection, the possibility that people who are already active or prefer to use and walk/cycle for transport purposes select to live in areas with good access to public transport (Cao, Mokhtarian, & Handy, 2009). Only one article explicitly controlled for self-selection (Schoner & Cao, 2014). Though LRT had no significant effect on walking after controlling for demographics, travel attitudes, and residential preferences, the models found significant effects of built environment characteristics on walking. Three mention this as a limitation of their study (Appleyard et al., 2019; Ewing & Hamidi, 2014; MacDonald et al., 2010) and another three state that natural experiments can reduce concerns about self-selection as the people surveyed chose to live in their neighbourhoods before there was an LRT station (Hong et al., 2016; Huang et al., 2017; Spears et al., 2017).

Discussion

This systematic review summarizes and assesses research on the associations between LRT, an increasingly popular public transport investment, and physical activity. The physical activity outcomes most studied were walking behaviour, cycling behaviour, walking and cycling behaviour combined, and MVPA.

Discussion of key findings

A moderate certainty of evidence found a positive association between LRT and walking behaviour. Studies found that LRT were generally associated with a 7–40% increase in walking with up to a 151% increase reported in one study. A strength of this work is the high proportion of natural experiment studies, as well as studies that calculated walking behaviour using devices. However, the studies included often report different types of walking behaviour (e.g. total walking, walking in an area, utilitarian walking, etc.), which makes direct comparison difficult. Given this, urban planners can be relatively confident that LRT investments will result in increased walking behaviour. It is also recommended that future research should carefully consider which types of walking behaviour they are measuring in their studies and aim to recruit bigger samples to solidify the evidence for policy makers.

Three studies examined the distance walked to LRT stations, two of which found that people walk – or are willing to walk – further to access LRT than buses. More studies on this relationship are needed. If future studies replicate this finding, the longer distances walked to LRT can be used to advocate for this form of public transport.

The certainty of evidence for a mixed relationship between LRT and MVPA is low; findings were often not statistically significant. These less robust results may be because LRT primarily encourages physical activity through walking to/from stations, a form of physical activity generally considered as lighter intensity. This may explain why overall physical activity decreased in Hirsch et al. (2018)'s meta-analysis while transport-related physical activity increased. As Hirsch et al. (2018) note, the "ActivityStat hypothesis", when people change their usual activity patterns to compensate for increased active travel, may also exist. Further, some studies found that baseline MVPA was associated with MVPA after LRT construction suggesting that those who are already physically active prior to the construction of LRT are likely to be more active post-construction – it is not clear if there was a shift amongst the population who are previously inactive. Though these studies were natural experiments, self-selection may confound the results (i.e. those who are more active to begin to live in neighbourhoods that are the sites of LRT investments). It is important to note that switching from private vehicle to LRT may result in other health, economic, and societal benefits, even if it does not result in an increase in MVPA.

Very low evidence exists for the relationship between LRT and cycling behaviour. Few use bicycles as a station access/egress mode in cities with low overall cycling rates (Pucher & Buehler, 2009; Ravensbergen, Buliung, Mendonca, & Garg, 2018). Perhaps the studies included did not consistently capture a significant relationship between LRT and cycling behaviour because cycling participation is already low prior to LRT construction. Alternatively, cyclists may replace some of their cycling trips after an LRT is built or

extended. Future work should examine whether stations in areas with higher cycling rates might exhibit a stronger relationship between physical activity from cycling and LRT use and/or proximity. The integration of cycling with public transit in cities dominated by the private automobile has been discussed as both a challenge and a great opportunity (Bachand-Marleau, Larsen, & El-Geneidy, 2011; Krizek & Stonebraker, 2010; Ravensbergen et al., 2018). Challenges include the lack of built environment characteristics that support cycling such as bicycle lanes, high quality parking, or even station design that accommodates bicycles (e.g. elevators with space for bicycles, a train car devoted to bicycles, etc.) (Ravensbergen et al., 2018). Perhaps changes in cycling behaviour take longer to develop than walking behaviour as the neighbourhood and station changes that might be required to enable cycling behaviour may take several years to develop. Given that cycling is a healthy and environmentally friendly travel mode that extends station catchment areas when compared to walking and reduces parking costs when compared to driving (Ravensbergen et al., 2018), the very low certainty of evidence identified herein should not discourage planners from designing LRT stations to encourage cycling. Future studies that carefully examine the connection of LRT stations to the existing cycling network and available bicycle parking are needed to better understand the rationale behind low cycling rates.

A further three studies examined walking and/or cycling behaviours across different LRT stations. All found some evidence that walking behaviour varies across LRT networks. Though this literature is limited, the results highlight the importance of considering the built environment and design features at LRT stations. In other words, it is possible that not all LRT or LRT stops are created equal when it comes to encouraging physical activity. Past research has examined how station design influences, amongst other things, access to jobs (Lahoorpoor & Levinson, 2020) and accessibility for people living with disabilities (Unsworth, So, Chua, Gudimetla, & Naweed, 2019). More research on how LRT station design influences physical activity (for instance, see: Loutzenheiser, 1997) is recommended and urban planners and policy makers are encouraged to consider how the built environment and station design influence physical activity as they plan LRT.

Strengths and weaknesses of the literature

Amongst the papers included in this review, strengths include a large number of studies using a natural experiment design to evaluate the introduction of a new LRT or LRT extension. Amongst cross-sectional studies, controlling for confounders, or comparing experimental groups to control groups was also common; however, only one study explicitly controlled for neighbourhood self-selection. Further, the use of both self-reported and device-measured physical activity data was complimentary providing both support for perceived behaviour, as well as movement intensity (Colley, Butler, Garriguet, Prince, & Roberts, 2018).

In terms of limitations, only one study included herein assessed physical activity at more than one time period after construction (Spears et al., 2017). Given that the association between LRT and physical activity was stronger after 6 months than after 18 months, there is a need for longitudinal research with repeated measure designs. Further, few studies reported measures of variation, including confidence

intervals, for physical activity outcome increases, a best practice in the field of public health. Finally, few studies explicitly examined whether the physical activity outcomes associated with LRT are distributed equally, and no studies examine how these outcomes affect children, youth, or older adults. Therefore, more research that examines the equity implications of LRT on physical activity among different age groups is needed.

Several research protocols for the evaluation of LRT on physical activity were identified. Though protocols were excluded from the review, many show promise for informing the gaps identified in this review, such as whether the impact of LRTs are distributed equitably (e.g. Durand et al., 2016; Roberts, Hu, Saksvig, Brachman, & Durand, 2018), as well as explicitly measuring the impacts of LRT on physical activity amongst youth (Roberts et al., 2018).

Conclusion

Taken together, this paper expands on past reviews examining the associations between public transport and health and Hirsch et al. (2018)'s meta-analysis on the relationship between rapid transit and physical activity. Given the heterogeneity in the reporting of the data, a comprehensive review of the association between LRT and physical activity is presented rather than a meta-analysis. A moderate certainty of evidence for a positive association between LRT and walking is identified. Further, low certainty of evidence for the mixed associations with MVPA and very low certainty of evidence and a lack of statistically significant association between LRT and cycling, and walking and cycling behaviour combined was identified.

Therefore, practitioners can be relatively confident that LRT investments will lead to changes in walking behaviour but should be cautious in assuming they will result in cycling or MVPA rates. There is a continued need for research examining the relationship between LRT and cycling behaviour, as well as experimental evidence with repeated measures that control for self-selection bias. Practitioners may also need to incorporate more considerations for cycling while planning for LRT by ensuring stations are accessible by safe bicycle infrastructure and include secure bicycle parking. Research examining whether the physical activity outcomes of LRT are distributed equitably, including how they impact different individuals and populations, such as children or older adults, as well as how physical activity varies across LRT stations, would also contribute to the literature.

Finally, while this paper provides practitioners with the certainty of the evidence between LRT and different types of physical activity, future work can compliment this review by replicating this analysis for different types of public transport or in different contexts. For instance, a literature review of LRT with a global scope could parse apart not only how LRTs impact different physical activity outcomes, but how these vary across countries, regions, cities, and potentially even neighbourhood typologies. This would allow for a comparison between the physical activity benefits associated with LRT in car-oriented contexts compared to contexts more conducive to active travel. Further, research comparing the impact of LRT on physical activity to that of other modes could help policy makers decide which public transport investments result in the greatest benefits.

Note

1. Because none of the outcome variables met the criteria to be upgraded, upgrading evaluation criteria are omitted from Appendix 4.

Acknowledgements

The authors would like to thank James DeWeese for his help conceptualizing the search strategy and contributing to the screening of papers and David Greene, the McGill librarian who helped develop the search strategy.

Disclosure statement

No potential conflict of interest was reported by the author(s).

Funding

This research was funded by The Canadian Institutes of Health Research (CIHR) and The Natural Sciences and Engineering Research Council of Canada (NSERC) Collaborative Health Research Projects (CHRP) Program (Project numbers CIHR CPG-170602 and CPG-170602 X-253156, NSERC CHRPJ 549576-20). The content and views expressed in this article are those of the authors and do not necessarily reflect those of the Government of Canada.

ORCID

Léa Ravensbergen  <http://orcid.org/0000-0003-3259-6673>

Rania Wasfi  <http://orcid.org/0000-0002-8605-1540>

Stephanie A. Prince  <http://orcid.org/0000-0001-6729-5649>

Gregory Butler  <http://orcid.org/0000-0002-7536-2044>

Yan Kestens  <http://orcid.org/0000-0003-2619-5750>

Ahmed El-Geneidy  <http://orcid.org/0000-0002-0942-4016>

References

- Appleyard, B., Frost, A., & Allen, C. (2019). Are all transit stations equal and equitable? Calculating sustainability, livability, health, & equity performance of smart growth & transit-oriented-development (TOD). *Journal of Transport & Health*, 14. doi:[10.1016/j.jth.2019.100584](https://doi.org/10.1016/j.jth.2019.100584)
- Bachand-Marleau, J., Larsen, J., & El-Geneidy, A. (2011). Much-anticipated marriage of cycling and transit: How will it work? *Transportation Research Record*, 2247(1), 109–117. doi:[10.3141/2247-13](https://doi.org/10.3141/2247-13)
- Balshem, H., Helfand, M., Schünemann, H. J., Oxman, A. D., Kunz, R., Brozek, J., ... Guyatt, G. H. (2011). GRADE guidelines: 3. Rating the quality of evidence. *Journal of Clinical Epidemiology*, 64(4), 401–406. doi:<https://doi.org/10.1016/j.jclinepi.2010.07.015>
- Barbaric, M., & Alizadeh, T. (2017). *Gold coast light rail performance-based public private partnerships: An effective delivery model?* Paper presented at the 8th State of Australian Cities National Conference, Adelaide, Australia. Retrieved from <https://apo.org.au/node/178496>
- Bopp, M., Gayah, V., & Campbell, M. (2015). Examining the link between public transit use and active commuting. *International Journal of Environmental Research and Public Health*, 12(4), 4256–4274.
- Brown, B., Smith, K., Jensen, W., & Tharp, D. (2017). Transit rider body mass index before and after completion of street light-rail line in Utah. *American Journal of Public Health*, 107(9), 1484–1486.
- Brown, B., Smith, K., Tharp, D., Werner, C., Tribby, C., Miller, H., & Jensen, W. (2016). A complete street intervention for walking to transit, nontransit walking, and bicycling: A quasi-experimental

- demonstration of increased use. *Journal of Physical Activity & Health*, 13(11), 1210–1219. doi:10.1123/jpah.2016-0066
- Brown, B., & Werner, C. (2007). A new rail stop – Tracking moderate physical activity bouts and ridership. *American Journal of Preventive Medicine*, 33(4), 306–309. doi:10.1016/j.amepre.2007.06.002
- Brown, B., Werner, C., Smith, K., Tribby, C., Miller, H., Jensen, W., & Tharp, D. (2016). Environmental, behavioral, and psychological predictors of transit ridership: Evidence from a community intervention. *Journal of Environmental Psychology*, 46, 188–196. doi:10.1016/j.jenvp.2016.04.010
- Cao, X., Mokhtarian, P., & Handy, S. (2009). Examining the impacts of residential self-selection on travel behaviour: A focus on empirical findings. *Transport Reviews*, 29(3), 359–395. doi:10.1080/01441640802539195
- Carlson, S., Watson, K., Paul, P., Schmid, T., & Fulton, A. (2017). Understanding the demographic differences in neighborhood walking supports. *Journal of Physical Activity and Health*, 14(4), 253–264. doi:10.1123/jpah.2016-0273
- Colley, R., Butler, B., Garriguet, D., Prince, S., & Roberts, K. (2018). *Comparison of self-reported and accelerometer-measured physical activity in Canadian adults* (Vol. 29, pp. 3–15). Statistics, Ottawa, Canada.
- Crist, K., Brondeel, R., Tuz-Zahra, F., Reuter, C., Sallis, J., Pratt, M., & Schipperijn, J. (2021). Correlates of active commuting, transport physical activity, and light rail use in a university setting. *Journal of Transport & Health*, 20. doi:10.1016/j.jth.2020.100978
- Daniel, J. (2012). Choosing the size of the sample. In *Sampling essentials: Practical guidelines for making sampling choices*, Johnnie Daniel (ed), Thousand Oaks, CA: Sage.
- Durand, C., Oluyomi, A., Gabriel, K., Salvo, D., Sener, I., Hoelscher, D., ... Kohl III, H. (2016). The effect of light rail transit on physical activity: Design and methods of the travel-related activity in neighborhoods study. *Frontiers in Public Health*, 4. doi:10.3389/fpubh.2016.00103
- Eady, J., & Burtt, D. (2019). *Walking and transport in Melbourne suburbs*.
- Ewing, R., & Hamidi, S. (2014). Longitudinal analysis of transit's land use multiplier in Portland (OR). *Journal of the American Planning Association*, 80(2), 123–137. doi:10.1080/01944363.2014.949506
- Guyatt, G., Oxman, A., Kunz, R., Brozek, J., Alonso-Coello, P., Rind, D., ... Vist, G. (2011). GRADE guidelines 6. Rating the quality of evidence—imprecision. *Journal of Clinical Epidemiology*, 64(12), 1283–1293.
- Hampton Roads Transit. (2015). *Final Report from the before-and-after study of the tide light rail project*.
- Hess, D. B., Ray, P., & Attard, N. (2014). *Connections beyond campus: An evaluation of the Niagara frontier transportation authority – University at Buffalo pilot transit pass program*.
- Hirsch, J., DeVries, D., Brauer, M., Frank, L., & Winters, M. (2018). Impact of new rapid transit on physical activity: A meta-analysis. *Preventive Medicine Reports*, 10, 184–190. doi:10.1016/j.pmedr.2018.03.008
- Hong, A., Boarnet, M., & Houston, D. (2016). New light rail transit and active travel: A longitudinal study. *Transportation Research Part a-Policy and Practice*, 92, 131–144. doi:10.1016/j.tra.2016.07.005
- Huang, R., Moudon, A., Zhou, C., Stewart, O., & Saelens, B. (2017). Light rail leads to more walking around station areas. *Journal of Transport & Health*, 6, 201–208. doi:10.1016/j.jth.2017.02.002
- Johnson, M. (2019). *Light rail? Heavy rail? Subway? Rail transit modes fall on a continuum*. Retrieved from <https://ggwash.org/view/71583/light-rail-heavy-rail-subway-rail-transit-modes-fall-on-a-continuum>
- Krizek, K., & Stonebraker, E. (2010). Bicycling and transit: A marriage unrealized. *Transportation Research Record*, 2144(1), 161–167. doi:10.3141/2144-18
- Kumar Maghelal, P. (2007). *Healthy transportation – healthy communities: Developing objective measures of built-environment using GIS and testing significance of pedestrian variables on walking to transit* (Doctor of Philosophy, 3281115). Ann Arbor, MI: Texas A&M University, ProQuest Information and Learning.
- Lachapelle, U., & Noland, R. (2012). Does the commute mode affect the frequency of walking behavior? The public transit link. *Transport Policy*, 21, 26–36.

- Lachapelle, U., & Pinto, D. (2016). Longer or more frequent walks: Examining the relationship between transit use and active transportation in Canada. *Journal of Transport & Health*, 3(2), 173–180.
- Lahoorpoor, B., & Levinson, D. (2020). Catchment if you can: The effect of station entrance and exit locations on accessibility. *Journal of Transport Geography*, 82, 102556. doi:[10.1016/j.jtrangeo.2019.102556](https://doi.org/10.1016/j.jtrangeo.2019.102556)
- Langlois, M., Wasfi, R. A., Ross, N. A., & El-Geneidy, A. M. (2016). Can transit-oriented developments help achieve the recommended weekly level of physical activity? *Journal of Transport & Health*, 3 (2), 181–190. doi:[10.1016/j.jth.2016.02.006](https://doi.org/10.1016/j.jth.2016.02.006)
- Loutzenheiser, D. (1997). Pedestrian access to transit: Model of walk trips and their design and urban form determinants around bay area rapid transit stations. *Transportation Research Record*, 1604(1), 40–49. doi:[10.3141/1604-06](https://doi.org/10.3141/1604-06)
- MacDonald, J., Stokes, R., Cohen, D., Kofner, A., & Ridgeway, G. (2010). The effect of light rail transit on body mass index and physical activity. *American Journal of Preventive Medicine*, 39(2), 105–112. doi:[10.1016/j.amepre.2010.03.016](https://doi.org/10.1016/j.amepre.2010.03.016)
- Malouff, D. (2015). *How to tell the difference between streetcars and light rail*. Retrieved from <https://ggwash.org/view/36980/how-to-tell-the-difference-between-streetcars-and-light-rail>
- McAslan, D. (2018). *Walking, transit use, and urban morphology in walkable urban neighborhoods: An examination of behaviors and attitudes in Seattle* (Doctor of Philosophy, 10902999). Ann Arbor, MI: University of Michigan ProQuest.
- Miller, H., Tribby, C., Brown, B., Smith, K., Werner, C., Wolf, J., ... Oliveira, M. (2015). Public transit generates new physical activity: Evidence from individual GPS and accelerometer data before and after light rail construction in a neighborhood of Salt Lake City, Utah, USA. *Health & Place*, 36, 8–17. doi:[10.1016/j.healthplace.2015.08.005](https://doi.org/10.1016/j.healthplace.2015.08.005)
- Murad, M. H., Mustafa, R. A., Schünemann, H. J., Sultan, S., & Santesso, N. (2017). Rating the certainty in evidence in the absence of a single estimate of effect. *Evidence-based Medicine*, 22(3), 85–87. doi:[10.1136/ebmed-2017-110668](https://doi.org/10.1136/ebmed-2017-110668)
- Noland, R., Ozbay, K., DiPetrillo, S., & Iyer, S. (2014). *Measuring benefits of transit oriented development*.
- O'Sullivan, S. (1995). *Walking distances to and from light rail transit stations in Calgary* (Master of Science in Civil Engineering). University of Calgary National Library Canada.
- Pucher, J., & Buehler, R. (2009). Integrating bicycling and public transport in North America. *Journal of Public Transportation*, 12(3), 79–104. doi:<http://doi.org/10.5038/2375-0901.12.3.5>
- Ravensbergen, L., Buliung, R., Mendonca, M., & Garg, N. (2018). Biking to ride: Investigating the challenges and barriers of integrating cycling with regional rail transit. *Transportation Research Record*, 2672(8), 374–383. doi:[10.1177/0361198118777080](https://doi.org/10.1177/0361198118777080)
- Rissel, C., Curac, N., Greenaway, M., & Bauman, A. (2012). Physical activity associated with public transport use – A review and modelling of potential benefits. *International Journal of Environmental Research and Public Health*, 9(7), 2454–2478. doi:[10.3390/ijerph9072454](https://doi.org/10.3390/ijerph9072454)
- Roberts, J., Hu, M., Saksvig, B., Brachman, M., & Durand, C. (2018). Examining the influence of a new light rail line on the health of a demographically diverse and understudied population within the Washington, DC metropolitan area: A protocol for a natural experiment study. *International Journal of Environmental Research and Public Health*, 15(2). doi:[10.3390/ijerph15020333](https://doi.org/10.3390/ijerph15020333)
- Schoner, J., & Cao, X. (2014). Walking for purpose and pleasure influences of light rail, built environment, and residential self-selection on pedestrian travel. *Transportation Research Record*, 2464, 67–76. doi:[10.3141/2464-09](https://doi.org/10.3141/2464-09)
- Sener, I., Lee, R., & Elgart, Z. (2016). Potential health implications and health cost reductions of transit-induced physical activity. *Journal of Transport & Health*, 3(2), 133–140. doi:[10.1016/j.jth.2016.02.002](https://doi.org/10.1016/j.jth.2016.02.002)
- Sinclair, O. (2019). Big year ahead for North American LRT. *Railway Age*. Retrieved from <https://www.railwayage.com/passenger/light-rail/big-year-ahead-for-north-american-lrt/>
- Spears, S., Boarnet, M. G., & Houston, D. (2017). Driving reduction after the introduction of light rail transit: Evidence from an experimental-control group evaluation of the Los Angeles Expo Line. *Urban Studies*, 54(12), 2780–2799. doi:[10.1177/0042098016657261](https://doi.org/10.1177/0042098016657261)

- The Transport Politic. (2021). *Route miles for US and Canada rail systems*. Retrieved from <https://www.thetransportpolitic.com/databook/route-miles-for-us-and-canada-rail-systems/>
- Unsworth, C., So, M., Chua, J., Gudimetla, P., & Naweed, A. (2019). A systematic review of public transport accessibility for people using mobility devices. *Disability and Rehabilitation*, 1–15. doi:10.1080/09638288.2019.1697382
- van Soest, D., Tight, M. R., & Rogers, C. (2020). Exploring the distances people walk to access public transport. *Transport Reviews*, 40(2), 160–182. doi:10.1080/01441647.2019.1575491
- Xiao, C., Goryakin, Y., & Cecchini, M. (2019). Physical activity levels and new public transit: A systematic review and meta-analysis. *American Journal of Preventive Medicine*, 56(3), 464–473. doi:10.1016/j.amepre.2018.10.022
- Zhang, Z., Liu, J., Li, X., & Khattak, A. (2021). Do larger sample sizes increase the reliability of traffic incident duration models? A case study of east Tennessee incidents. *Transportation Research Record*, 1–16. doi:10.1177/0361198121992063

Appendices

Appendix 1. Database search strategy

Database	Limits	Search terms	
		LRT	Physical activity
Web of Science (n = 2266)	TS (topic)	"Light Rail Transit" OR "Light Rail" OR LRT OR "light metro" OR "Urban rail" OR subway OR metro	physical activit* OR exercise* OR walk* OR bik* OR cycl* OR bicycl* OR LMPA OR MVPA OR METS OR metabolic rate*
TRID (n = 184)	Title search	Light Rail Transit OR Light Rail OR LRT OR light metro OR urban rail OR subway OR metro	physical activit* OR exercis* OR walk* OR bik* OR bicycl* OR cycl* OR LMPA OR MVPA OR METS OR metabolic rate*
Scopus (n = 2835)	Title, abstract, keywords	"Light Rail Transit" OR "Light Rail" OR LRT OR "light metro" OR "Urban rail" OR subway OR metro	"physical activit*" OR exercise* OR walk* OR bik* OR cycl* OR bicycl* OR LMPA OR MVPA OR METS OR metabolic rate*
Medline (n = 430)	Title, abstract, keyword heading	Light Rail Transit OR Light Rail OR LRT OR light metro OR Urban rail OR subway OR metro	physical activit* OR exercise* OR walk* OR bik* OR cycl* OR bicycl* OR LMPA OR MVPA OR METS OR metabolic rate*
SportsDiscus (n = 151)	Title, abstract, keywords	Light Rail Transit OR Light Rail OR LRT OR light metro OR Urban rail OR subway OR metro	physical activit* OR exercise* OR walk* OR bik* OR cycl* OR bicycl* OR LMPA OR MVPA OR METS OR metabolic rate*

Note: Web of Science (2266), TRID (184), Scopus (2835), Medline (430), SportsDiscus (151).

Appendix 2. Cities with LRT included in the grey literature search

Country	City
Australia	City of Adelaide City of Melbourne City of Gold Coast City of Newcastle City of Sydney City of Canberra
New Zealand U.S.A.	City of Auckland (LRT under construction) City of Baltimore City of Boston City of Buffalo City of Charlotte City of Cleveland City of Dallas City of Denver City of Houston Jersey City City of Los Angeles City of Minneapolis City of Newark City of Norfolk City of Philadelphia City of Phoenix City of Pittsburgh City of Portland City of Sacramento City of St-Louis Salt Lake City City of San Diego City of San Francisco City of San Jose/Santa Clara City of Seattle City of Calgary
Canada	

(Continued)

Continued.

Country	City
	City of Edmonton
	City of Kitchener-Waterloo
	City of Ottawa

Appendix 3. Effective Public Health Practice Project (EPHPP) quality assessment tool for quantitative studies

Criteria type	Criteria	Evaluation
Components rating		
Selection bias	<p>(Q1) Are the individuals selected to participate in the study likely to be representative of the target population?</p> <p>1 Very likely 2 Somewhat likely 3 Not likely 4 Can't tell</p> <p>(Q2) What percentage of selected individuals agreed to participate?</p> <p>1 80–100% agreement 2 40–79% agreement 3 less than 40% agreement 4 Not applicable 5 Can't tell</p>	<p>Strong = sample's socio-demographic characteristics are compared to the general population (e.g. through the census) and little discrepancy is identified OR 60% or more agreed to participate</p> <p>Moderate = sample's socio-demographic characteristics are compared to the general population (e.g. through the census) and some discrepancy is identified OR use multiple data sources, some of which are highly representative (e.g. the census) OR representative sampling strategy</p> <p>Weak = no comparison to general population and very small sample (under 100)</p>
Study design	<p>Indicate the study design</p> <p>Was the study described as randomized? No/Yes</p> <p>If Yes, was the method of randomization described? No/Yes</p> <p>If Yes, was the method appropriate? No/Yes</p>	<p>Strong = natural experiment design</p> <p>Moderate = randomized cross-sectional design</p> <p>Weak = cross-sectional design with no randomization</p>
Confounders	<p>(Q1) Were there important differences between groups prior to the intervention?</p> <p>1 Yes 2 No 3 Can't tell</p> <p>The following are examples of confounders: Race, Sex, Marital status/family, Age, SES (income or class), Education, Health status, Pre-intervention score on outcome measure</p> <p>(Q2) If yes, indicate the percentage of relevant confounders that were controlled (either in the design (e.g. stratification, matching) or analysis)?</p> <p>1 80–100% (most) 2 60–79% (some) 3 Less than 60% (few or none) 4 Can't tell</p>	<p>Strong = compared groups and found minimal differences</p> <p>Moderate = compared groups and found some differences OR controlled well for confounders</p> <p>Weak = did not compare groups OR did not control well for confounders</p>
Blinding	<p>(Q1) Was (were) the outcome assessor(s) aware of the intervention or exposure status of participants?</p> <p>1 Yes 2 No 3 Can't tell</p> <p>(Q2) Were the study participants aware of the research question?</p> <p>1 Yes 2 No 3 Can't tell</p>	<p>Strong = authors state the participants were not aware of research question during data collection</p> <p>Moderate = unclear whether participants were not aware of research question during data collection</p> <p>Weak = paper states participants were aware of research question during data collection</p>

(Continued)

Continued.

Criteria type	Criteria	Evaluation
Data collection methods	(Q1) Were data collection tools shown to be valid? 1 Yes 2 No 3 Can't tell (Q2) Were data collection tools shown to be reliable? 1 Yes 2 No 3 Can't tell	Strong = data collection tools result in measured data (accelerometry) Moderate = self-reported, but vigorous, survey Weak = self-reported through poorly described survey OR data originate from different sources
Withdrawals and drop-outs	(Q1) Were withdrawals and drop-outs reported in terms of numbers and/or reasons per group? 1 Yes 2 No 3 Can't tell 4 Not applicable (i.e. one time surveys or interviews) (Q2) Indicate the percentage of participants completing the study. (If the percentage differs by groups, record the lowest) 1 80–100% 2 60–79% 3 less than 60% 4 Can't tell 5 Not applicable (i.e. retrospective case-control)	Strong = withdrawal and drop-outs reported (numbers and reasons), minimal differences identified between withdrawals and sample Moderate = withdrawal and drop-outs reported (numbers and reasons) with some differences identified OR 60% or more participants completed the study Weak = withdrawals and dropouts not reported. Less than 60% completed the study Not applicable = all cross-sectional research designs
Intervention integrity	(Q1) What percentage of participants received the allocated intervention or exposure of interest? 1 80–100% 2 60–79% 3 less than 60% 4 Can't tell (Q2) Was the consistency of the intervention measured? 1 Yes 2 No 3 Can't tell (Q3) Is it likely that subjects received an unintended intervention (contamination or co-intervention) that may influence the results? 4 Yes 5 No 6 Can't tell	Strong = consistency of intervention measured (e.g. distance to LRT) AND no reason to believe participants received an unintended intervention beyond regular daily life Moderate = no reason to believe participants received an unintended intervention beyond regular daily life Weak = reason to believe participants received an unintended intervention beyond regular daily life Not applicable = all cross-sectional research designs
Analyses	(Q1) What statistical methods are used? (Q2) Are the statistical methods appropriate for the study design? 1 Yes 2 No 3 Can't tell	Strong = utilizes statistical modelling Moderate = utilizes descriptive statistics Weak = presents results with no statistical analysis or unclear analysis
Global rating List all component ratings		Strong = no "weak" ratings on any component Moderate = one "weak" rating on a component Weak = two or more "weak" ratings on any components

Appendix 4. GRADE evaluation of evidence criteria

Domain	Assessment	Score	Criteria
Risk of bias	No or little risk of bias	0	At least 75% of peer-reviewed papers scored moderate or strong on the following Risk of Bias Assessment indicators: Confounders, Blinding, Data Collection Methods, Withdrawals and Drop-Outs, and Intervention Integrity
	Moderate risk of bias	-0.5	50–74% of peer-reviewed papers scored moderate or strong on any of the following Risk of Bias Assessment indicators: Confounders, Blinding, Data Collection Methods, Withdrawals and Drop-Outs, and Intervention Integrity
	High risk of bias	-1	Less than 50% of peer-reviewed papers scored moderate or strong on any of Risk of Bias Assessment indicators: Confounders, Blinding, Data Collection Methods, Withdrawals and Drop-Outs, and Intervention Integrity
Inconsistency	No or little inconsistency	0	At least 75% of studies identify the same expected direction of effect (e.g. positive association between physical activity and LRT). In the case of only one study per outcome, inconsistency was not relevant
	High inconsistency	-1	Less than 75% of studies identify the same expected direction of effect (e.g. positive association between physical activity and LRT)
Indirectness	Little concern	0	At least 75% of studies score strong or moderate on Selection Bias (i.e. how generalizable the results are Q1) in Risk of Bias Assessment
	High concern	-1	>75% of studies score strong on Selection Bias in Risk of Bias Assessment
Imprecision	No or little imprecision	0	Optimal information size at least 1000 ^a for natural experiment OR 6500 ^b for cross-sectional
	High imprecision	-1	Optimal information size under 1000 for natural experiment OR 6500 for cross-sectional
Publication bias	No or little publication bias	0	Sample size is larger than 400 ^c in 75% or more of studies that show expected associations
	Potential publication bias	-0.5	Sample size is less than 400 in 75% or more of studies
	High publication bias	-1	In at least 75% of studies, sample size is smaller than 400 Only cross-sectional studies show positive associations
Original level of certainty	Grade		Final level of certainty
High	0.5–1 point deducted		Moderate
	1.5–2 points deducted		Low
	2.5+ points deducted		Very low
	No points deducted		High
Low	Any points deducted		Very low
	No points deducted and eligible for an upgrade		Moderate

^aBased on recommendations in Guyatt et al. (2011).

^bThreshold deemed reliable in the models developed by Zhang, Liu, Li, and Khattak (2021).

^cThreshold derived from the “rule of thumb” for survey research, single topic in Daniel (2012).